

Speculative generalization of black hole uniqueness to higher dimensions

Barak Kol

School of Natural Sciences

Institute for Advanced Study

Einstein Drive, Princeton NJ 08540, USA

email barak@sns.ias.edu

A straightforward generalization of the celebrated uniqueness theorem to dimensions greater than four was recently found to fail in two pure gravity cases - the 5d rotating black ring and the black string on $\mathbf{R}^{3,1} \times \mathbf{S}^1$. Two amendments are suggested here (without proof) in order to rectify the situation. The first is that in addition to specifying the mass and angular momentum (and gauge charges) one needs to specify the horizon topology as well. Secondly, the theorem may survive if applied exclusively to stable solutions. Note that the latter is at odds with the proposed stable but non-uniform string.

PACS: 04.50.+h; 04.70.Bw hep-th/0208056

I. INTRODUCTION

The celebrated uniqueness theorem states that 4d black holes with regular horizons are determined uniquely by their mass M and angular momentum J , $|J|/M^2 \leq 1$ in pure gravity and by their electric (and possibly magnetic) charge when a Maxwell field is present, with important contributions by Israel, Carter, Hawking, Robinson, Mazur, Bunting and Masood Ul Alam and others. See for example [1–3] and for a thorough reference list see the reviews [4–7] and references therein.

Recently two cases were studied in which a straightforward generalization of black hole uniqueness to more than four dimensions fails. The first is the black-string to black-hole transition which takes place on any compactification background the simplest being $\mathbf{R}^{3,1} \times \mathbf{S}^1$ background – 3+1 extended dimensions and a fifth compact dimension of radius L , which is denoted here by the z coordinate. Black objects with Schwarzschild radius much smaller than L are expected to closely resemble a 5d Schwarzschild black hole (BH) with a nearly round $\mathbf{S}^3$ horizon topology, while very massive black objects, will have a horizon size much larger than L and will be extended over the z axis wrapping it completely. Such objects will have $\mathbf{S}^2 \times \mathbf{S}^1$ horizon topology and will be referred to as “black strings”. One type of black string is the 4d Schwarzschild solution with the z coordinate added in as a spectator and it will be called here a “uniform string”, while another type of solution may be z dependent, namely, a “non-uniform string”. Gregory and Laflamme discovered an onset of tachyonic instability [8,9] when the uniform black string becomes so thin that its Schwarzschild radius is of the same order as L . Following the claimed existence of a stable non-uniform solution [10] Gubser performed a higher order analysis of the Gregory-Laflamme (GL) critical point and found that the non-uniform solution which must emerge there (as the GL mode is marginally tachyonic) is unstable and its mass is larger than the critical GL mass [11].

Later an analysis of the phase diagram of the whole process, supplemented by a local demonstration of topology change near the cone over $\mathbf{S}^2 \times \mathbf{S}^2$ produced the following phase diagram (figure 1) [12,13]. In this diagram it was not possible to accommodate the stable non-uniform string without creating some other serious problem. The violation of uniqueness is apparent for a range of masses above the GL critical mass, where three solutions are present: the uniform string, the 5d black hole and an unstable (unlike the previous two) and non-uniform string. Closer to the merger point probably there are even more phases but with more unstable modes as well.




FIG. 1. A suggested phase diagram for the black-hole black-string transition [12]. μ is the dimensionless parameter $G_5 M/L^2$, and λ is an order parameter (measure of non-uniformity). In the range $\mu_{GL} \leq \mu \leq \mu_{\text{merger}}$ (at least) three solutions coexist. Solid lines are stable phases while dashed ones are unstable. The solutions to the left of the merger point are black strings while those to the right are black holes. Dotted lines denote transitions - a first order transition at μ_1 and tachyonic decay from the two other points. The full phase diagram should include further refinements in the area of the merger transition.

The second example is the 5 dimensional rotating black object (in uncompactified pure gravity). Here there are two kinds of solutions, a black hole with horizon topology $\mathbf{S}^3$ [14] and the recently discovered rotating ring with horizon $\mathbf{S}^2 \times \mathbf{S}^1$ [15]. For given mass M the black hole exists for an angular momentum range $|J| \leq J_2$ for some J_2 , while the ring exists for $J \geq J_1$ for some $J_1 < J_2$. As can be seen in figure (2) uniqueness is violated in the range $J_1 < J < J_2$ (as noted by the authors) where three solutions exist: two rings and one black hole. The stability/ instability of either of these solutions is not known at this time. Moreover, in 5d there are two parameters of angular momentum a_1, a_2 and this picture emerged for $a_2 = 0$ but it is not known yet how it generalizes to $a_2 \neq 0$.




FIG. 2. Rotating solutions in 5d. $(27\pi/32G)J^2/M^3$ as a function of some parameter ν . The dashed line is the rotating black hole of [14] while the solid line is the newly discovered black ring [15]. In the range $J_1 \leq J \leq J_2$ marked by the dotted lines three solutions coexist, denoted I, II and III from right to left. Reproduced with permission from [15].

Following [15] uniqueness in higher dimensions was reexamined and the straightforward generalization was found to hold for the static ($J = 0$) uncompactified case and also when some scalar fields are included [16–19]. Other relatively recent related works include [20–22].

Here a different approach will be taken, and rather than trying to map the set of cases where straightforward uniqueness holds and where it does not, I will speculate

on an appropriate generalization of uniqueness that may hold without exception, hoping such a generalization exists since considering the fundamental importance of the uniqueness theorem in 4d it would be a shame if it simply disappeared in higher dimensions.

The uniqueness theorem can be thought as being made out of two steps - the counting of parameters of black holes (“no hair”) and the impossibility of bifurcations or disconnected solution families. The parameter counting is the fundamental physical property, related to the amount of required boundary conditions, and it is not violated in higher dimensions. Actually, once the counting is done for small black holes with small angular momenta, it is bound to hold always including for black holes in the presence of a compactification or a cosmological constant. Thus we are left to examine bifurcations and disconnected components, which rely on a different principle for proof - the positive energy theorems.

Guided by the phenomenology, namely the two cases described above, it is seen that bifurcations do occur, and two criteria suggest themselves to me. The first is that the *horizon topology* should be added to the list of (continuous) characterizing parameters such as mass etc. Yet in both cases this is not enough to uniquely specify a solution. Thus the second criterion of *stability* is suggested. Actually this is better thought to be a raised question, namely does uniqueness hold if solutions (with identical horizon topologies) are limited to be stable? This condition is physically attractive since unstable solutions cannot be attained anyway (due to noise).

The other possibility is that there is no generalization of uniqueness to higher dimensions and phase diagrams with several phases are common without any simple constraints on the number or type of phases.

II. HORIZON TOPOLOGY

In 4 dimensions the only possible topology for a horizon is $\mathbf{S}^2$ (forgetting the light-like direction $\simeq \mathbf{R}$) [23]. However, the examples above show that in higher dimensions the horizon topology is not unique. Since any discussion of uniqueness of time independent solutions starts with a dimensional reduction along Killing vectors and analysis of the boundary conditions, where the boundary is composed of asymptotic infinity and the event horizon, it is clear that the horizon topology must be specified as one of these boundary conditions.

One may ask what are the possible horizon topologies in higher dimensions. From the examples it is seen that in 5d both $\mathbf{S}^3$ and $\mathbf{S}^2 \times \mathbf{S}^1$ are possible. By considering toroidal compactifications a generalized black string in d dimensions could have *any* $\mathbf{S}^k \times \mathbf{T}^{d-2-k}$ topology where $\mathbf{T}^n \simeq (\mathbf{S}^1)^n$ is the n -dimensional torus, and $k \geq 2$.

More generally we may consider compactification over an arbitrary Ricci flat manifold, and consider the changes in topology of a small black hole as its mass is being in-

creased. The possible topologies are not known, other than the obvious restriction that they should be cobordant with the topology at asymptotic infinity (and the cobordism type may be constrained in certain theories). A possible guide for topologies could be the change in topology of the equipotential lines of a (scalar) point source on the compactification manifold as one goes away from the source. It would be interesting to find a simple characterization of these topologies, if one exists.

III. STABILITY

In both of the cases studied here horizon topology is not enough to specify a solution in the parameter range of coexistence. Thus I am led to wonder whether adding a stability requirement would do. In the case of the black-hole black-string transition there are two overlapping black-string solutions but the non-uniform one is unstable [11] and so the “generalized uniqueness” holds. Note that *if* the latter is indeed correct that would constitute another argument against the existence of the stable non-uniform string of [10], at least in the range of parameters where the uniform string is stable.

For the rotating ring there are again three solutions in the overlap region. Suppose we start with the black hole solution at $J = J_1$, trace out the solution line (for fixed M), and denote by region I the black hole solutions with J going from J_1 to J_2 , by region II the black ring solutions going from J_2 to J_1 and finally by region III the black ring going from J_1 to J_2 . Since a stability analysis is not available at this time, it is possible to provide a *concrete test* for the stability criterion – namely that the black ring solutions in region II (on the right in figure 2) are unstable (actually it would be enough if regions II and III have instability domains which together cover the region $J_1 \leq J \leq J_2$).

But why is there any reason to believe that the stability criterion would be correct? The main tool so far in proving uniqueness has been the positive energy theorems which may be thought to be a strong generalization of the Green identity for the Laplace equation, which is used to prove uniqueness for the boundary value problem (either Dirichlet or Neumann). However, this tool makes no reference to stability and so it must fail in cases where several solutions exist. A possible direction is to consider the phase transitions in the system. If there were two stable phases with the same horizon topology then when one of them would decay to the other, it could do so without a topology change and so presumably without going through an explosive singularity [12,13]. It is conceivable that such a mild phase transition is forbidden by some “censorship”.

Acknowledgments

It is a pleasure to thank G. Gibbons for comments.

Work supported by DOE under grant no. DE-FG02-90ER40542, and by a Raymond and Beverly Sackler Fellowship.

-
- [1] W. Israel, “Event Horizons In Static Vacuum Space-Times,” *Phys. Rev.* **164**, 1776 (1967).
 - [2] B. Carter, “Axisymmetric Black Hole Has Only Two Degrees of Freedom,” *Phys. Rev. Lett.* **26**, 331 (1971).
 - [3] D. C. Robinson, “Uniqueness of the Kerr Black Hole,” *Phys. Rev. Lett.* **34**, 905 (1975).
 - [4] P. O. Mazur, “Black hole uniqueness theorems,” *arXiv:hep-th/0101012*.
 - [5] M. Heusler, “Stationary black holes: Uniqueness and beyond,” *Living Rev. Rel.* **1**, 6 (1998).
 - [6] B. Carter, “Mathematical Foundations of the Theory of Relativistic Stellar and Black Hole Configurations,” in B. Carter and J. B. Hartle eds., *Gravitation in Astrophysics*, 63-122, (Plenum Press, New York, 1987).
 - [7] P. T. Chrusciel, “‘No Hair’ Theorems: Folklore, Conjectures, Results,” *Contemp. Math.* **170**, 23 (1994) [*arXiv:gr-qc/9402032*].
 - [8] R. Gregory and R. Laflamme, “Black Strings And P-Branes Are Unstable,” *Phys. Rev. Lett.* **70**, 2837 (1993) [*arXiv:hep-th/9301052*].
 - [9] R. Gregory and R. Laflamme, “The Instability of charged black strings and p-branes,” *Nucl. Phys. B* **428**, 399 (1994) [*arXiv:hep-th/9404071*].
 - [10] G. T. Horowitz and K. Maeda, “Fate of the black string instability,” *Phys. Rev. Lett.* **87**, 131301 (2001) [*arXiv:hep-th/0105111*].
 - [11] S. S. Gubser, “On non-uniform black branes,” *arXiv:hep-th/0110193*.
 - [12] B. Kol, “Topology change in general relativity and the black-hole black-string transition,” *arXiv:hep-th/0206220*.
 - [13] B. Kol, “Explosive black hole fission and fusion in large extra dimensions,” *arXiv:hep-ph/0207037*.
 - [14] R. C. Myers and M. J. Perry, “Black Holes In Higher Dimensional Space-Times,” *Annals Phys.* **172**, 304 (1986).
 - [15] R. Emparan and H. S. Reall, “A rotating black ring in five dimensions,” *Phys. Rev. Lett.* **88**, 101101 (2002) [*arXiv:hep-th/0110260*].
 - [16] S. Hwang “A Rigidity Theorem for Ricci Flat Metrics,” *Geometriae Dedicata* **71**, 5 (1998).
 - [17] G. W. Gibbons, D. Ida and T. Shiromizu, “Uniqueness and non-uniqueness of static vacuum black holes in higher dimensions,” *arXiv:gr-qc/0203004*,
 - [18] G. W. Gibbons, D. Ida and T. Shiromizu, “Uniqueness and non-uniqueness of static black holes in higher dimensions,” *and Phys. Rev. Lett.* **89**, 041101 (2002) [*arXiv:hep-th/0206049*],

- [19] G. W. Gibbons, D. Ida and T. Shiromizu, “Uniqueness of (dilatonic) charged black holes and black p-branes in higher dimensions,” and arXiv:hep-th/0206136.
- [20] M. Rogatko, “Uniqueness theorem for static dilaton $U(1)^*N$ black holes,” *Class. Quant. Grav.* **19**, 875 (2002) [arXiv:hep-th/0202132].
- [21] M. Mars, “Uniqueness properties of the Kerr metric,” *Class. Quant. Grav.* **17**, 3353 (2000) [arXiv:gr-qc/0004018].
- [22] C. G. Wells, “Extending the black hole uniqueness theorems. I: Accelerating black holes: The Ernst solution and C-metric,” arXiv:gr-qc/9808044.
- [23] S. W. Hawking, “The event horizon,” in *Black Holes*, eds. C. De Witt and B. S. De Witt, pp. 5-34, New York, Gordon and Breach.